

# Time reversal methods in unknown medium and inverse problems

Kenrick Bingham  
Yaroslav Kurylev  
Matti Lassas  
Samuli Siltanen

January 4, 2007

## Abstract

A novel method to solve inverse problems for the wave equation is introduced. The method is a combination of the boundary control method and an iterative time reversal scheme, leading to adaptive imaging of coefficient functions of the wave equation using focusing waves in unknown medium. The approach is computationally effective since the iteration lets the medium do most of the processing of the data.

The iterative time reversal scheme also gives an algorithm for constructing boundary controls for which the corresponding final values are as close as possible to the final values of a given wave in a part of the domain, and as close as possible to zero elsewhere. The algorithm does not assume that the coefficients of the wave equation are known.

**Keywords:** Inverse problems, wave equation, control, time reversal.

**AMS classification:** 35R30, 93B05.

## 1. Introduction

We present a novel inversion method for the wave equation. Suppose that we can send waves from the boundary into an unknown body with spatially varying wave speed  $c(x)$ . Using a combination of the boundary control (BC) method and an iterative time reversal scheme, we show how to focus waves near a point  $x_0$  inside the medium and simultaneously recover  $c(x_0)$  if  $c$  is

isotropic. In the anisotropic case we can reconstruct  $c(x)$  up to a change of coordinates.

In the classical BC method material parameters are reconstructed using hyperbolic techniques, see [1, 9, 10, 11, 25, 26, 27]. Straightforward numerical implementation of the BC method is computationally demanding. Inspired by Isaacson's iterative measurement scheme [22] for electrical impedance tomography (EIT), we combine the BC method with an adaptive time reversal iteration that lets the medium do most processing of the data.

Traditional time reversal methods record waves, invert them in time, and send them back into the medium. If the recorded signal originated from point sources or was reflected from small scatterers in the medium, the time reversed waves focus at the source or scatterer points. This is useful for time reversal mirrors in communication technologies and medical therapies. For early references on time reversal (using ultrasound in air), see the seminal works of Fink [19, 18, 17]. For a microlocal discussion of time reversal, see Bardos [3, 4], and for another mathematical treatment see Klivanov [29].

Time reversal in known background medium with random fluctuations has also been extensively studied [5, 6, 7, 8, 13, 14] and applied to medical imaging, non-destructive testing and underwater acoustics. These methods are outside the scope of this paper.

Iterative time reversal has been used to find best measurements for inverse problems. By “best” we here mean “optimal for detecting the presence of an object”. This *distinguishability problem* has been studied for fixed-frequency problems in EIT [22] and acoustic scattering [34]. The connection between optimal measurements and iterative time-reversal experiments was pointed out in [34, 35, 36]. For the wave equation, the best measurement problem has been studied in [16], where it has been shown that the optimal incident field for probing a half space can be found by an iterative process involving time reversal mirrors.

Let us describe our hybrid method in a simple case. Take a compact set  $M \subset \mathbb{R}^3$  with smooth boundary  $\partial M$ , and let  $c(x)$  be a scalar-valued wave speed in  $M$ . Consider the wave equation

$$\begin{aligned} u_{tt} - c(x)^2 \Delta u &= 0 && \text{in } M \times \mathbb{R}_+, \\ u|_{t=0} &= 0, \\ u_t|_{t=0} &= 0, \\ -c(x)^{-2} \partial_n u &= f(x, t) && \text{in } \partial M \times \mathbb{R}_+, \end{aligned} \tag{1}$$

where  $\partial_n$  denotes the Euclidean normal derivative. We denote by  $u^f = u^f(x, t)$  the solution of (1) corresponding to a given boundary source term  $f$ .

Take  $T > 0$  larger than the diameter of  $M$  in travel time metric, so that any point inside  $M$  can be reached by waves sent from the boundary in time less than  $T$ . We consider imposing a boundary source  $f$  on the set  $\partial M \times (0, 2T)$  and measuring the boundary value of the resulting wave  $u^f$  during the time  $0 < t < 2T$ . All such boundary measurements constitute the Neumann-to-Dirichlet map

$$\Lambda_{2T} : f \mapsto u^f|_{\partial M \times (0, 2T)},$$

that models a variety of physical measurements [28]. Instead of the full map we only need here the values  $\Lambda_{2T}f$  for a specific collection of sources  $f$  given by an iterative process.

We need three special operators on the function space  $L^2(\partial M \times [0, 2T])$ . First, the *time reversal operator*

$$Rf(x, t) = f(x, 2T - t),$$

second, the *time filter*

$$Jf(x, t) = \frac{1}{2} \int_0^{\min(2T-t, t)} f(x, s) ds,$$

and finally the *restriction operator*

$$Pf(x, t) = \chi_B(x, t)u(x, t),$$

where  $\chi_B$  is the characteristic function of a set  $B \subset \partial M \times [0, 2T]$ .

We define the *processed time reversal iteration*

$$\begin{aligned} F &:= \frac{1}{\omega} P(R\Lambda_{2T}RJ - J\Lambda_{2T})f, \\ a_n &:= \Lambda_{2T}(h_n), \quad b_n := \Lambda_{2T}(RJh_n), \\ h_{n+1} &:= (1 - \frac{\alpha}{\omega})h_n - \frac{1}{\omega}(PRb_n - PJa_n) + F, \end{aligned} \tag{2}$$

where  $f \in L^2(\partial M \times [0, 2T])$  and  $\alpha, \omega > 0$  are parameters. Starting with  $h_0 = 0$  the iteration (2) converges to a limit  $h(\alpha) = \lim_{n \rightarrow \infty} h_n$ . In the case  $B = \partial M \times [T - s, T]$  we have

$$\lim_{\alpha \rightarrow 0} u^{h(\alpha)}(x, T) = (1 - \chi_N(x))u^f(x, T) \quad \text{in } L^2(M), \tag{3}$$

where  $N \subset M$  is the set of points  $x \in M$  with travel time distance to  $\partial M$  at most  $s$ . In other words,  $N$  is the maximal region where the waves sent from the boundary can propagate in time  $s$ .

Let  $y \in \partial M$  and denote by  $\gamma_{y,\nu}$  the geodesic (in travel time metric) starting from  $y$  in the normal direction and parametrised by arclength. Take a source  $f$  concentrated near  $(y, T-s) \in \partial M \times \mathbb{R}_+$  and vanishing for  $t \leq T-s$ . Then  $u^f(x, T)$  is supported in the union of the set  $N$  and a neighbourhood of the point  $x_0 = \gamma_{y,\nu}(s)$ , and the limit (3) vanishes outside a neighbourhood of  $x_0$ . This way our iteration produces a wave localized near  $x_0$  at time  $T$ .

Below we introduce an alternative construction of focusing waves for general sources  $f$  by applying the processed time reversal iteration with two different sets  $B$  and using the difference of the resulting boundary values.

We show also that the limits (3) can be used to determine the travel time distance between an arbitrary point  $x = \gamma_{y,\nu}(s) \in M$  and any boundary point  $z \in \partial M$ . It follows that the processed time reversal iteration gives an algorithm for determining the wave speed in the medium. Furthermore, given a source  $f$ , our iteration allows us to find a wave  $u^h(x, t)$  having at time  $t = T_1$  approximately the value  $\chi_N(x)u^f(x, T_2)$  for any  $T_2$  without knowing the material parameters of the medium. Analogous methods without the multiplier  $\chi_N(x)$  have been developed before in [23, 31].

The paper is organised as follows. In Section 2 we formulate our main results. In Section 3 we prove the convergence of the processed time reversal iteration. Section 4 is devoted to the analysis of the focusing properties of the waves, and in Section 5 we show how to reconstruct the metric using the boundary distance function. In the last section we discuss our method in the case of noisy measurements.

## 2. Definitions and main results

Let us consider the closure  $M \subset \mathbb{R}^m$ ,  $m \geq 1$ , of an open smooth set, or a (non-compact or compact) complete Riemannian manifold  $(M, g)$  of dimension  $m$  with a non-empty boundary. For simplicity, we assume that the boundary  $\partial M$  is compact. Let  $u$  solve the wave equation

$$\begin{aligned} u_{tt}(x, t) + Au(x, t) &= 0 \quad \text{in} \quad M \times \mathbb{R}_+, \\ u|_{t=0} &= 0, \quad u_t|_{t=0} = 0, \\ B_{\nu,\eta}u|_{\partial M \times \mathbb{R}_+} &= f. \end{aligned} \tag{4}$$

Here,  $f \in L^2(\partial M \times \mathbb{R}_+)$  is a real valued function,  $A$  is a formally self-adjoint elliptic partial differential operator of the form (in local coordinates in the case when  $M$  is a manifold)

$$Av = - \sum_{j,k=1}^m \mu(x)^{-1} |g(x)|^{-\frac{1}{2}} \frac{\partial}{\partial x^j} \left( \mu(x) |g(x)|^{\frac{1}{2}} g^{jk}(x) \frac{\partial v}{\partial x^k}(x) \right) + q(x)v(x) \tag{5}$$

where  $[g^{jk}]$  is a smooth real positive definite matrix,  $c_0 I \leq [g^{jk}(x)] \leq c_1 I$ ,  $c_0, c_1 > 0$ ,  $|g| = \det([g_{jk}])$ , where  $[g_{jk}]$  is the inverse matrix of  $[g^{jk}]$ , and  $\mu(x) \geq c_2 > 0$  and  $q(x)$  are smooth real valued functions. Also,

$$B_{\nu, \eta} v = -\partial_\nu v + \eta v$$

where  $\eta : \partial M \rightarrow \mathbb{R}$  is a smooth bounded function and

$$\partial_\nu v = \sum_{j,k=1}^m \mu(x) g^{jk}(x) \nu_k \frac{\partial}{\partial x^j} v(x),$$

where  $\nu(x) = (\nu_1, \nu_2, \dots, \nu_m)$  is the interior co-normal vector field of  $\partial M$  normalised so that  $\sum_{j,k=1}^m g^{jk} \nu_j \nu_k = 1$ . A particular example is the operator

$$A_0 = -c^2(x) \Delta + q(x) \quad (6)$$

for which  $\partial_\nu v = c(x)^{-m+1} \partial_n v$ , where  $\partial_n v$  is the Euclidean normal derivative of  $v$ .

We denote the solutions of (4) by

$$u^f(x, t) = u(x, t).$$

For the initial boundary value problem (4) we define the response operator (or non-stationary Robin-to-Dirichlet map)  $\Lambda$  by setting

$$\Lambda f = u^f|_{\partial M \times \mathbb{R}_+}. \quad (7)$$

We also consider the finite time response operator  $\Lambda_T$  corresponding to time  $T > 0$ ,

$$\Lambda_T f = u^f|_{\partial M \times (0, T)}. \quad (8)$$

By [39], the map  $\Lambda_T : L^2(\partial M \times (0, T)) \rightarrow H^{1/3}(\partial M \times (0, T))$  is bounded, where  $H^s(\partial M \times (0, T))$  denotes the Sobolev space on  $\partial M \times (0, T)$ . Below we consider  $\Lambda_T$  as a bounded operator that maps  $L^2(\partial M \times (0, T))$  to itself.

We also need some other notation. The matrix  $g_{jk}(x)$ , that is, the inverse of the matrix  $g^{jk}(x)$ , is a Riemannian metric in  $M$  that is called the travel time metric. The reason for this is the waves propagate with speed one with respect to the metric  $ds^2 = \sum_{j,k} g_{jk}(x) dx^j dx^k$ . We denote by  $d(x, y)$  the distance function corresponding to this metric. For the wave equation we define the space  $L^2(M, dV_\mu)$  with inner product

$$\langle u, v \rangle_{L^2(M, dV_\mu)} = \int_M u(x) v(x) dV_\mu(x),$$

where  $dV_\mu = \mu(x)|g(x)|^{1/2}dx^1dx^2 \dots dx^m$ .

For  $t > 0$  and  $\Gamma \subset \partial M$ , let

$$M(\Gamma, t) = \{x \in M : d(x, \Gamma) \leq t\}, \quad (9)$$

be the domain of influence of  $\Gamma$  at time  $t$ .

We denote  $L^2(B) = \{f \in L^2(\partial M \times \mathbb{R}_+) : \text{supp}(f) \subset B\}$ ,  $B \subset \partial M \times \mathbb{R}_+$ , identifying functions and their zero continuations. When  $\Gamma \subset \partial M$  is open set and  $f \in L^2(\Gamma \times \mathbb{R}_+)$ , it is well known (see e.g. [21]) that the wave  $u^f(t) = u^f(\cdot, t)$  is supported in the domain  $M(\Gamma, t)$ ,

$$u^f(t) \in L^2(M(\Gamma, t)) = \{v \in L^2(M) : \text{supp}(v) \subset M(\Gamma, t)\}.$$

### 2.1. Results on convergence of processed time reversal iteration.

Our objective is to find a boundary source  $f$  such that the wave  $u^f(x, T)$  is localized in a neighbourhood of a single point  $x_0$ . Note that in this paper we do not focus the pair  $(u(x, T), u_t(x, T))$ , but only the value of the wave,  $u(x, T)$ .

Let  $0 < T_0 < T$ ,  $\Gamma \subset \partial M$  be open set and  $f \in L^2(\Gamma \times \mathbb{R}_+)$ . Our first aim is to construct boundary values  $h(\alpha) \in L^2(\partial M \times \mathbb{R}_+)$ ,  $\alpha > 0$  such that

$$\lim_{\alpha \rightarrow 0} u^{h(\alpha)}(T) = \chi_{M(\partial M, T_0)} u^f(T) \quad (10)$$

in  $L^2(M)$ . Here  $\chi_X(x)$  is the characteristic function of the set  $X$ . Then

$$\lim_{\alpha \rightarrow 0} u^{f-h(\alpha)}(T) = (1 - \chi_{M(\partial M, T_0)}) u^f(T)$$

is supported in  $M(\Gamma, T) \setminus M(\partial M, T_0)$  (see Figure 1). When  $T_0 \rightarrow T$  and  $\Gamma_j \rightarrow \{z\}$  then under suitable assumptions discussed later the set  $M(\Gamma_j, T) \setminus M(\partial M, T_0)$  goes to a single point. Thus for  $f \in L^2(\Gamma_j \times \mathbb{R}_+)$ , the waves  $(1 - \chi_{M(\partial M, T_0)}) u^f(T)$  localize to a small neighbourhood of a single point. For later use we will consider (10) below in a more general setting.


Next we explain a procedure to find boundary values  $h(\alpha)$ . Denote

$$\begin{aligned} R_{2T} f(x, t) &= f(x, 2T - t), \\ J_{2T} h(x, t) &= \int_{[0, 2T]} J(s, t) h(x, s) ds \end{aligned} \quad (11)$$

where  $J(s, t) = \frac{1}{2} \chi_L(s, t)$ ,

$$L = \{(s, t) \in \mathbb{R}_+ \times \mathbb{R}_+ : t + s \leq 2T, \ s > t\}. \quad (12)$$

Figure 1: The set  $M(\Gamma, T) \setminus M(\partial M, T_0)$ . Our goal is to find waves sent from the boundary that focus into this area.



We call  $R = R_{2T}$  the *time reversal map* and  $J = J_{2T}$  the *time filter*. We denote by  $P = P_B : L^2(\partial M \times [0, 2T]) \rightarrow L^2(\partial M \times [0, 2T])$  the multiplication operator

$$P_B f(x, t) = \chi_B(x, t) f(x, t),$$

where  $B = \bigcup_{j=1}^J (\Gamma_j \times [T - T_j, T])$ ,  $\Gamma_j \subset \partial M$  are open sets and  $0 \leq T_j \leq T$ . Next, we consider  $\Lambda_{2T}$ ,  $R_{2T}$ , and  $J_{2T}$  as operators from  $L^2(\partial M \times [0, 2T])$  to itself.

Let  $\alpha \in (0, 1)$  and  $\omega > 0$  be a sufficiently large constant. We define  $a_n, b_n \in L^2(\partial M \times [0, 2T])$  and  $h_n = h_n(\alpha) \in L^2(\partial M \times [0, 2T])$  by the iteration

$$\begin{aligned} a_n &:= \Lambda_{2T}(h_n), \quad b_n := \Lambda_{2T}(R J h_n), \\ h_{n+1} &:= (1 - \frac{\alpha}{\omega}) h_n - \frac{1}{\omega} (P R b_n - P J a_n) + F, \end{aligned} \tag{13}$$

with  $a_0 = 0$ ,  $b_0 = 0$ ,  $h_0 = 0$ , and

$$F = \frac{1}{\omega} P (R \Lambda_{2T} R J - J \Lambda_{2T}) f.$$

We say that this iteration is the *processed time reversal iteration on time-interval  $[0, 2T]$  with projector  $P$*  and starting point  $f \in L^2(\partial M \times [0, 2T])$ .

Here  $a_n$  corresponds to the “iterated measurement”,  $b_n$  to the “time filtered and time-reversed measurement” and  $h_{n+1}$  to post-processing of  $h_n$ ,  $a_n$  and  $b_n$  using time reversal and time filtering.

**Theorem 1** *Let  $T > 0$ . Assume we are given  $\partial M$  and the response operator  $\Lambda_{2T}$ . Let  $\Gamma_j \subset \partial M$ ,  $j = 1, \dots, J$  be non-empty open sets,  $0 \leq T_j \leq T$ , and  $B = \bigcup_{j=1}^J (\Gamma_j \times [T - T_j, T])$ . Let  $f \in L^2(\partial M \times \mathbb{R}_+)$  and let for  $\omega$  large enough and  $\alpha \in (0, 1)$  functions  $h_n = h_n(\alpha)$  be defined by the processed time reversal iteration (13) with projector  $P_B$  and starting point  $f_0 = f|_{\partial M \times (0, 2T)}$ . These functions converge in  $L^2(\partial M \times \mathbb{R}_+)$ ,*

$$h(\alpha) = \lim_{n \rightarrow \infty} h_n(\alpha)$$

*and the limits satisfy*

$$\lim_{\alpha \rightarrow 0} u^{h(\alpha)}(x, T) = \chi_N(x) u^f(x, T)$$

*in  $L^2(M)$ , where  $N = \bigcup_{j=1}^J M(\Gamma_j, T_j) \subset M$ .*

This theorem is proven later in Section 3. Before that we consider its consequences.

**2.2. Results on focusing of the waves** Let us consider a geodesic  $\gamma_{x,\xi}$  in  $(M, g)$  parametrised along arc length where  $\gamma_{x,\xi}(0) = x$ ,  $\dot{\gamma}_{x,\xi}(0) = \xi$  with  $\|\xi\|_g = 1$ . Let  $\nu = \nu(z)$ ,  $z \in \partial M$  be the unit interior normal vector of  $\partial M$  in  $(M, g)$ . There is a critical value  $\tau(z) \in (0, \infty]$ , such that for  $t < \tau(z)$  the geodesic  $\gamma_{z,\nu}([0, t])$  is the unique shortest geodesic, and for  $t > \tau(z)$  it is no longer a shortest geodesic from its endpoint  $\gamma_{z,\nu}(t)$  to  $\partial M$ .

We say that  $\Gamma_j \rightarrow \{\widehat{z}\}$  if  $\Gamma_{j+1} \subset \Gamma_j$  and  $\bigcap_{j=1}^{\infty} \Gamma_j = \{\widehat{z}\}$ .

Theorem 1 yields the following results telling that we can produce focusing waves, that is, wave that focus to a single point.

**Corollary 2** *Let  $\widehat{z} \in \partial M$  and  $0 < T_0 < \widehat{T} < T$ . Let  $\widehat{x} = \gamma_{\widehat{z},\nu}(\widehat{T})$  and  $\Gamma_j \subset \partial M$ ,  $j \in \mathbb{Z}_+$  be open neighbourhoods of  $\widehat{z} \in \partial M$  such that  $\Gamma_j \rightarrow \{\widehat{z}\}$  when  $j \rightarrow \infty$ .*


*Let  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$ . Let  $h_n(\alpha; T_0, j)$  be the functions obtained from the processed time reversal iteration (13) with projector  $P_B$ ,  $B = (\Gamma_j \times [T - \widehat{T}, T]) \cup (\partial M \times [T - T_0, T])$  and starting point  $f_0 = f|_{\partial M \times (0, 2T)}$ . Similarly, let  $h'_n(\alpha; T_0, j)$  be the functions obtained from the processed time reversal iteration (13) with projector  $P_{B'}$ ,  $B' = \partial M \times [T - T_0, T]$  and starting point  $f_0$ . Let*

$$\widetilde{h}_n(\alpha; T_0, j) = f_0 - h_n(\alpha; T_0, j) + h'_n(\alpha; T_0, j).$$

*If  $\widehat{T} < \tau(\widehat{z})$  then*

$$\lim_{T_0 \rightarrow \widehat{T}} \lim_{j \rightarrow \infty} \lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{(\widehat{T} - T_0)^{(m+1)/2}} u^{\widetilde{h}_n(\alpha; T_0, j)}(T) = C_0(\widehat{x}) u^f(\widehat{x}, T) \delta_{\widehat{x}}(x) \quad (14)$$

Figure 2: Subsets of  $M$  used in the proof of Corollary 2.



in  $\mathcal{D}'(M)$  where  $C_0(\hat{x}) > 0$  does not depend on  $f$ . If  $\hat{T} > \tau(\hat{z})$  the limit (14) is zero.

In Figure 2, the set  $M(\Gamma_j, \hat{T}) \setminus M(\partial M, T_0)$  tends to the point  $\hat{x}$  when  $j \rightarrow \infty$  and  $T_0 \rightarrow \hat{T}$ . This turns out to be essential in the proof of Corollary 2.

Above,  $\delta_{\hat{x}}$  is Dirac's delta distribution on  $(M, g)$  such that

$$\int_M \delta_{\hat{x}}(x) \phi(x) dV_\mu = \phi(\hat{x}), \quad \phi \in C_0^\infty(M).$$

**2.3. Inner products.** Later, we show that that using  $f, h \in L^2(\partial M \times [0, 2T])$  and the boundary measurements  $\Lambda_{2T}$  we may compute via an explicit integral the inner product

$$\int_M u^f(x, T) u^h(x, T) dV_\mu(x) = \int_{\partial M \times [0, 2T]} (Kf)(x, t) h(x, t) dS_g(x) dt, \quad (15)$$

where  $dS_g$  is the Riemannian surface volume of  $\partial M$  and

$$K = K_{2T} := R_{2T} \Lambda_{2T} R_{2T} J_{2T} - J_{2T} \Lambda_{2T} \quad (16)$$

so that terms (15) can be found using measurement operator  $\Lambda_{2T}$  and simple basic operations like time reversal  $R_{2T}$  and the time filter operator  $J_{2T}$ .

In the above formula the Riemannian surface volume of  $\partial M$  can be computed using the intrinsic metric of  $\partial M$ , which is determined by the map  $\Lambda_{2T}$ . Indeed, it follows from Tataru's unique continuation principle (see [38, 40], see also Theorem 4 below) that the Schwartz kernel of  $\Lambda_{2T}$  is supported in the set  $E = \{(x, t, x', t') \in (\partial M \times [0, 2T])^2 : t - t' \geq d(x, x')\}$  and that the boundary  $\partial E$  is in the support. The set  $\partial E$  determines the distances of points  $z, z' \in \partial M$  with respect to the metric of  $(M, g)$ , and thus also the distance with respect to the intrinsic metric of the boundary  $(\partial M, g_{\partial M})$ .

**2.4. Reconstruction of material parameter functions.** It is well known that the response map  $\Lambda$  can not generally determine coefficients  $\mu$  and  $g_{jk}$  because of two transformations discussed below.

First, we one can introduce a coordinate transformation, that is a diffeomorphism  $F : M \rightarrow M$  such that the boundary value  $F|_{\partial M}$  is the identity operator. Then the push forward metric  $\tilde{g} = F_*g$ , that is,

$$\tilde{g}_{jk}(y) = \sum_{p,q=1}^m \frac{\partial x^p}{\partial y^j} \frac{\partial x^q}{\partial y^k} g_{pq}(x), \quad y = F(x)$$

and the functions  $\tilde{\mu} = \mu \circ F^{-1}$ ,  $\tilde{q} = q \circ F^{-1}$ , and  $\tilde{\eta} = \eta$  determine the operator  $\tilde{A}$  of the form (5) such that response operators for  $A$  and  $\tilde{A}$  are the same (for this, see [25, 26, 27]).

Second, one can do the gauge transformation  $u(x, t) \rightarrow \kappa(x)u(x, t)$  where  $\kappa \in C^\infty(M)$  is strictly positive function. In this transformation the operator  $\tilde{A}$  is transforms to the operator  $\tilde{A}_\kappa$  defined be

$$\tilde{A}_\kappa w := \kappa \tilde{A} \left( \frac{1}{\kappa} w \right)$$

and the operator in boundary condition is transformed to  $B_{\nu, \hat{\eta}}$ , with

$$\hat{\eta} = \eta - \kappa^{-1} \partial_\nu \kappa.$$

Then the response operator  $\tilde{\Lambda}_\kappa$  of  $\tilde{A}_\kappa$  coincides with the response operator  $\Lambda$  of  $A$ .

By above transformations, making a gauge transformation with  $\kappa = \mu^{-1}$ , we come to operator  $\tilde{A}_\kappa$  having the same response operator as  $A$  and that can be represented in the form (5) with  $\mu(x) = 1$ . It turns out that this is the only source of non-uniqueness.

**Corollary 3** *Assume that  $\mu = 1$  and that we are given the boundary  $\partial M$  and the response operator  $\Lambda$ . Then using the the processed time reversal iteration we can find constructively the manifold  $(M, g)$  upto an isometry and on it the operator  $A$  uniquely.*

*Moreover, if  $M \subset \mathbb{R}^m$ ,  $m \geq 2$ , and  $A$  is of the form (6), given the set  $M$  and the response operator  $\Lambda$  we can determine  $c(x)$  and  $q(x)$  uniquely in  $M$ .*

We note that the unique determination of  $(M, g)$  and  $A$  is well known, see e.g. [11, 25, 26] and references therein. The novelty of Corollary 3 is that in the construction based on processed time reversal iteration the use of iterated measurement avoid many computationally demanding steps used required in [26]. An example of such step is a Gram-Schmidt orthogonalisation of basis of spaces  $L^2(\Gamma \times [t_1, t_2])$ ,  $\Gamma \subset \partial M$  with respect to an inner product determined by  $\Lambda_{2T}$ . This is a typical step used in traditional Boundary Control method that is very sensitive for measurement errors.

### 3. Proof of the convergence of the iteration

**3.1. Controllability results.** The seminal result implying controllability is Tataru's unique continuation result (see [38, 40])

**Theorem 4 (Tataru)** *Let  $u$  be a solution of wave equation*

$$u_{tt}(x, t) + Au(x, t) = 0.$$

*Assume that*

$$u|_{\Gamma \times (0, 2\tau)} = 0, \quad \partial_\nu u|_{\Gamma \times (0, 2\tau)} = 0 \quad (17)$$

*where  $\Gamma \subset \partial M$ ,  $\Gamma \neq \emptyset$  is open, and  $\tau > 0$ . Then,*

$$u(x, \tau) = 0, \quad \partial_t u(x, \tau) = 0 \quad \text{for } x \in M(\Gamma, \tau).$$

This result yields the following Tataru's controllability result, see e.g. [26] and references therein.

**Theorem 5** *Let  $\Gamma \subset \partial M$  be open and  $\tau > 0$ . Then the linear subspace*

$$\{u^f(\tau) \in L^2(M(\Gamma, \tau)) : f \in C_0^\infty(\Gamma \times [0, \tau])\}$$

*is dense in  $L^2(M(\Gamma, \tau))$ .*

**3.2. Inner products.** Let us recall the Blagovestchenskii identity:

**Lemma 6** *Let  $f, h \in L^2(\partial M \times [0, 2T])$ . Then*

$$\begin{aligned} & \int_M u^f(x, T) u^h(x, T) dV_\mu(x) = \\ & \int_{[0, 2T]^2} \int_{\partial M} J(t, s) [f(t)(\Lambda_{2T} h)(s) - (\Lambda_{2T} f)(t)h(s)] dS_g(x) dt ds, \end{aligned} \quad (18)$$

*where  $J(t, s) = \frac{1}{2} \chi_L(s, t)$ , see (12).*

The proof in slightly different context is given e.g. in [26].

**Proof.** Let  $w(t, s) = \int_M u^f(t) u^h(s) dV_\mu$ . Integrating by parts, we see that

$$\begin{aligned} & (\partial_t^2 - \partial_s^2)w(t, s) \\ &= - \int_M [Au^f(t)u^h(s) - u^f(t)Au^h(s)] dV_\mu(x) \\ &= - \int_{\partial M} [\partial_\nu u^f(t)u^h(s) - u^f(t)\partial_\nu u^h(s)] dS_g \\ &= \int_{\partial M} [(-\partial_\nu u^f(t) + \eta u^f(t))u^h(s) - u^f(t)(-\partial_\nu u^h(s) + \eta u^h(s))] dS_g \\ &= \int_{\partial M} [f(t)\Lambda_{2T}h(s) - \Lambda_{2T}f(t)h(s)] dS_g. \end{aligned} \quad (19)$$

Moreover,

$$w|_{t=0} = w|_{s=0} = 0, \quad \partial_t w|_{t=0} = \partial_s w|_{s=0} = 0.$$

Thus we can consider (19) as one dimensional wave equation with known right hand side and vanishing initial and boundary data. Solving this initial boundary value problem we see the claim.  $\square$

Consider now the operator  $K$  appearing in formula (15) in more detail. The Schwartz kernel of  $\Lambda_{2T}$  is the Dirichlet boundary value of the Green's function  $G(x, x', t - t')$  satisfying

$$\begin{aligned} (\partial_t^2 + A)G_{x',t'}(x, t) &= \delta_{x'}(x)\delta(t - t') \quad \text{in } M \times \mathbb{R}_+, \\ G_{x',t'}|_{t=0} &= 0, \quad \partial_t G_{x',t'}|_{t=0} = 0, \\ B_{\nu,\eta}G_{x',t'}|_{\partial M \times \mathbb{R}_+} &= 0, \end{aligned} \tag{20}$$

where  $G_{x',t'}(x, t) = G(x, x', t - t')$ . Then

$$G(x, x', t - t') = G(x', x, t - t')$$

and we see that

$$\Lambda_{2T}^* = R_{2T}\Lambda_{2T}R_{2T}$$

where  $R_{2T}f(x, t) = f(x, 2T - t)$  is the time reversal map.

Thus, as  $\Lambda_{2T}^* = R_{2T}\Lambda_{2T}R_{2T}$ , we can write result of Lemma 6 as

$$\int_M u^f(x, T)u^h(x, T) dV_\mu(x) = \int_{\partial M \times [0, 2T]} (Kf)(x, t) h(x, t) dS_g(x) dt \tag{21}$$

where  $K$  is defined in (16).

**3.3. Proof of Theorem 1.** To produce focusing waves, let us consider the problem

$$\min_{h \in L^2(B)} \|u^f(T) - u^h(T)\|_{L^2(M, dV_\mu)}^2 \tag{22}$$

where  $B = \bigcup_{j=1}^J (\Gamma_j \times [T - T_j, T])$ ,  $\Gamma_j \subset \partial M$  are open and  $0 \leq T_j \leq T$ .

As the solution of this minimisation problem does not always exist, we can regularise the problem and study

$$\min_{h \in L^2(\partial M \times [0, 2T])} F(h, \alpha) \tag{23}$$

where  $\alpha \in (0, 1)$  and

$$F(h, \alpha) = \langle K(Ph - f), Ph - f \rangle_{L^2(\partial M \times [0, 2T], dS_g)} + \alpha \|h\|_{L^2(\partial M \times [0, 2T], dS_g)}^2.$$

We recall that  $P = P_B$  is multiplication with the characteristic function of  $B$ , that is,  $(P_B h)(x, t) = \chi_B(x, t) \cdot h(x, t)$ .

By (21),

$$F(h, \alpha) = \|u^f(T) - u^{Ph}(T)\|_{L^2(M, dV_\mu)}^2 + \alpha \|h\|_{L^2(\partial M \times [0, 2T])}^2.$$

The minimisation problem (23) is equivalent to (22) when  $\alpha = 0$ .

**Lemma 7** *For given  $\alpha \in (0, 1)$  the problem (23) has a unique minimiser. Moreover, the minimiser is the unique solution of the equation*

$$(PKP + \alpha)h = PKf. \quad (24)$$

**Proof.** The minimisation problem is strictly convex, and the map  $h \mapsto u^h(T)$  is continuous  $L^2(\partial M \times [0, 2T]) \rightarrow H^{s_1}(M)$ ,  $s_1 < 5/6$  by [33], see also [32]. Now if  $h_j \rightarrow h$  weakly in  $L^2(\partial M \times [0, 2T])$  as  $j \rightarrow \infty$  then  $Ph_j \rightarrow Ph$  weakly in  $L^2(\partial M \times [0, 2T])$ . As the embedding  $H^{s_1}(M) \rightarrow L^2(M)$  is compact and the map  $h \mapsto u^h(T)$  is continuous  $L^2(\partial M \times [0, 2T]) \rightarrow H^{s_1}(M)$ , then  $u^{Ph_j}(T) \rightarrow u^{Ph}(T)$  in  $L^2(M)$ . Therefore,

$$\begin{aligned} & \lim_{j \rightarrow \infty} \langle K(Ph_j - f), Ph_j - f \rangle_{L^2(\partial M \times [0, 2T])} \\ &= \lim_{j \rightarrow \infty} \|u^{Ph_j}(T) - u^f(T)\|_{L^2(M, dV_\mu)}^2 = \|u^{Ph}(T) - u^f(T)\|_{L^2(M, dV_\mu)}^2. \end{aligned}$$

Let  $h_j$  be a sequence such that  $\lim_{j \rightarrow \infty} F(h_j, \alpha) = \inf_h F(h, \alpha)$ . As  $F(h, \alpha) \geq \alpha \|h\|_{L^2}^2$ , the sequence  $(h_j)$  is bounded. Since  $\|h\|_{L^2} \leq \liminf_{j \rightarrow \infty} \|h_j\|_{L^2}$ , this implies that  $F(h, \alpha) \leq \liminf_{j \rightarrow \infty} F(h_j, \alpha)$ .

Thus by choosing a subsequence of  $h_j$ , we can assume that  $h_j$  converge to  $h$  in the weak topology of  $L^2(\partial M \times [0, 2T])$ , and the limit  $h$  is a global minimiser of  $F(\cdot, \alpha)$ .

By computing the Fréchet derivative of the functional  $F(h, \alpha)$  in any direction  $\tilde{h} \in L^2(\partial M \times [0, 2T])$  at a minimiser  $h$ , we see that

$$\begin{aligned} 0 &= D_h(\langle K(Ph - f), Ph - f \rangle + \alpha \|h\|_{L^2}^2) \tilde{h} \\ &= \langle \tilde{h}, P^* K(Ph - f) \rangle + \langle \tilde{h}, P^* K^*(Ph - f) \rangle + 2\alpha \langle \tilde{h}, h \rangle \end{aligned}$$

for all  $\tilde{h}$ . Since  $K^* = K$  and  $P^* = P$ , this implies that  $h$  satisfies the equation (24). As  $K$  is non-negative,  $PKP + \alpha I \geq \alpha I$  so that equation (24) has a unique solution providing the minimiser  $h = h(\alpha)$ .  $\square$

Note that equation (24) also implies that

$$h(\alpha) \in \text{Ran}(P). \quad (25)$$

Next we want to solve equation (24) using iteration. To this end, let  $\omega \in \mathbb{R}_+$  be a constant such that

$$\omega > 2(1 + \|PKP\|).$$

Then the equation (24) can be written as

$$(I - S)h = \frac{1}{\omega} PKf, \quad \text{where} \quad S = I - \frac{\alpha + PKP}{\omega}. \quad (26)$$

Since  $PKP$  is non-negative and

$$\alpha I \leq \alpha I + PKP \leq \frac{\omega}{2} I,$$

we see that  $\|S\| < 1$ . Thus we can solve  $h$  using iteration: Let

$$F := \frac{1}{\omega} PKf = \frac{1}{\omega} P(J\Lambda_{2T} - R\Lambda_{2T}RJ)f,$$

$h_0 = 0$ , and consider the iteration

$$h_{n+1} = Sh_n + F, \quad n = 0, 1, \dots$$

As  $h_n = Ph_n$  by (25), we can write the iteration in the form (13), and  $\lim_{n \rightarrow \infty} h_n = h(\alpha)$  in  $L^2(\partial M \times [0, 2T])$ .

Applying this algorithm can find the minimisers  $h = h(\alpha) \in \text{Ran}(P)$  of problem (24). They converge by the following lemma:

**Lemma 8** *We have*

$$\lim_{\alpha \rightarrow 0} u^{h(\alpha)}(x, T) = \chi_N(x)u^f(x, T)$$

in  $L^2(M)$ , with  $N$  as in Theorem 1.

**Proof.** Clearly,

$$\begin{aligned} F(h, \alpha) &= F_0 + \|\chi_N(u^{Ph}(T) - u^f(T))\|_{L^2(M)}^2 + \alpha \|h\|_{L^2(\partial M \times [0, 2T])}^2, \\ F_0 &= \|(1 - \chi_N)u^f(T)\|_{L^2(M)}^2. \end{aligned} \quad (27)$$

Here the first term does not depend on  $h$ . By Theorem 5, for any  $\epsilon > 0$  there is a  $h_\epsilon \in L^2(B)$  such that

$$\|u^{Ph_\epsilon}(T) - \chi_N u^f(T)\|_{L^2(M)}^2 < \frac{\epsilon}{2}. \quad (28)$$

Thus we have

$$F(h_\epsilon, \alpha) = F_0 + \frac{\epsilon}{2} + \alpha \|h(\epsilon)\|_{L^2(\partial M \times [0, 2T])}^2.$$

This shows that if  $\alpha < \alpha(\epsilon)$ , where

$$\alpha(\epsilon) := \frac{\epsilon}{2\|h(\epsilon)\|_{L^2(\partial M \times [0, 2T])}^2},$$

then

$$F(h_\epsilon, \alpha) \leq F_0 + \epsilon.$$

Thus the minimiser  $h(\alpha)$  of  $F(h, \alpha)$  with  $\alpha < \alpha(\epsilon)$  satisfies

$$F(h(\alpha), \alpha) \leq F_0 + \epsilon.$$

As by (25),  $h(\alpha) = Ph(\alpha)$  so that  $h(\alpha) \in L^2(\Gamma \times [0, T])$ , we get from (28) that, for  $\alpha < \alpha(\epsilon)$ ,

$$\|\chi_N(u^{h(\alpha)}(T) - u^f(T))\|_{L^2(M)}^2 \leq \epsilon/2. \quad (29)$$

As  $\text{supp}(u^{h(\alpha)}(T)) \subset N$ , the claim follows.  $\square$

Theorem 1 follows from Lemmata 7 and 8.  $\square$

## 4. Proofs for the focusing of the waves

Next we prove Corollary 2.

**Proof of Corollary 2.** Let  $(z(x), s(x))$  be the boundary normal coordinates of  $x \in M$ , that is,  $s(x) = d(x, \partial M)$  and  $z(x)$  is the closest point of  $\partial M$  to  $x$  when such a point is unique. When a closest boundary point is not unique, the boundary normal coordinates are not defined.

We consider first the claim of the corollary in the case when  $\widehat{T} < \tau(\widehat{z})$ . Then the boundary normal coordinates near  $\widehat{x} = \gamma_{\widehat{z}, \nu}(\widehat{T})$  are well defined and the metric tensor in these coordinates has the form

$$g = \begin{pmatrix} 1 & 0 \\ 0 & [g_{\alpha\beta}(z, s)]_{\alpha, \beta=1}^{m-1} \end{pmatrix}, \quad (30)$$

where  $[g_{\alpha\beta}(z, s)] \in \mathbb{R}^{(m-1) \times (m-1)}$  is a smooth positive definite matrix valued function near  $(\widehat{z}, \widehat{T}) = (z(\widehat{x}), s(\widehat{x}))$ . Then there is a  $C_1 = C_1(\widehat{z}, \widehat{T}) > 0$  such that

$$C_1^{-1}I \leq [g_{\alpha\beta}(z, s)] \leq C_1I$$

near  $(\widehat{z}, \widehat{T})$ . Using this we see that there is a  $C_2 > 0$  such that

$$C_2^{-1} \leq \frac{\text{vol}_g(M(\{\widehat{z}\}, \widehat{T}) \setminus M(\partial M, T_0))}{(\widehat{T} - T_0)(\widehat{T}^2 - T_0^2)^{(m-1)/2}} \leq C_2.$$

As the solution  $u^f(x, t)$  is smooth for  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$ ,

$$\lim_{T_0 \rightarrow \widehat{T}} \frac{\chi_{M(\{\widehat{z}\}, \widehat{T}) \cup M(\partial M, T_0)} - \chi_{M(\partial M, T_0)}}{\text{vol}_g(M(\{\widehat{z}\}, \widehat{T}) \setminus M(\partial M, T_0))} u^f(T) = u^f(\widehat{x}, T) \delta_{\widehat{x}}$$

in  $\mathcal{D}'(M)$ , the claim follows in the case  $\widehat{T} < \tau(\widehat{z})$ .

When  $\widehat{T} > \tau(z)$ , the claim follows from the fact that for sufficiently small  $\widehat{T} - T_0 > 0$  the set  $M(\{\widehat{z}\}, \widehat{T}) \setminus M(\partial M, T_0)$  is empty.  $\square$

**Remark** The above proof of Corollary 2 also shows that in Corollary 2

$$C_0(x_0) = \lim_{T_0 \rightarrow \widehat{T}} \frac{(\widehat{T} - T_0)^{(m+1)/2}}{\text{vol}_g(M(\{\widehat{z}\}, \widehat{T}) \setminus M(\partial M, T_0))}. \quad (31)$$

## 5. Boundary distance functions and reconstruction of the metric

Here we present a method for finding the boundary distance functions using processed time reversal iteration.

Let  $z, y \in \partial M$ ,  $0 \leq T_1 \leq \tau(z)$  and  $T > d_{\partial M}(y, z) + T_1$ . Denote  $x = \gamma_{z, \nu}(T_1)$ . Next we give an algorithm that can be used to determine  $d(x, z)$ .

To this end, let  $\Gamma_j \subset \partial M$  and  $\Sigma_j \subset \partial M$  be neighbourhoods of  $z$  and  $y$ , respectively, such that  $\Gamma_j \rightarrow \{z\}$  and  $\Sigma_j \rightarrow \{y\}$  when  $j \rightarrow \infty$ .

Let  $\epsilon > 0$ ,  $\tau \in [0, T]$ , and

$$\begin{aligned} N_j^1 &= M(\Gamma_j, T_1) & B_j^1 &= \Gamma_j \times [T - T_1, T] \\ N_j^2 &= M(\Sigma_j, \tau) & B_j^2 &= \Sigma_j \times [T - \tau, T] \\ N_\epsilon^3 &= M(\partial M, T_1 - \epsilon) & B_\epsilon^3 &= \partial M \times [T - (T_1 - \epsilon), T]. \end{aligned}$$

**Lemma 9** *The distance  $d(x, z)$  is the infimum of all  $\tau \in [0, T]$  that satisfy the condition*

$$\begin{aligned} &\text{the set } I(j, \epsilon) = (N_j^1 \cap N_j^2) \setminus N_\epsilon^3 \text{ contains} \\ &\text{a non-empty open set for all } j \in \mathbb{Z}_+, \epsilon > 0. \end{aligned} \quad (32)$$




Figure 3: Sketch of the situation in Lemma 9 when  $I(j, \epsilon)$  is empty (left) and non-empty (right).

See Figure 3 for a sketch of the cases of empty and non-empty  $I(j, \epsilon)$ .

**Proof.** First consider what happens if  $d(x, y) < \tau$ . Since  $T_1 \leq \tau(z)$ , we see that then  $B(x, r) \cap (N_j^1 \setminus N_\epsilon^3)$  contains a non-empty open set for all  $r > 0$ , where  $B(x, r) \subset M$  is a ball of  $(M, g)$  with center  $x$  and radius  $r$ . When  $r < \tau - d(x, y)$ , we see that  $B(x, r) \subset N_j^2$ . Thus  $I(j, \epsilon)$  contains an open set and (32) is satisfied.

On other hand, if  $d(x, y) > \tau$ , let  $r = d(x, y) - \tau$ . When  $j \rightarrow \infty$  and  $\epsilon \rightarrow 0$ , we see using metric in the boundary normal coordinates (30) that  $N_j^1 \setminus N_\epsilon^3 \rightarrow \{x\}$  in the Hausdorff metric. Thus when  $j$  is large enough and  $\epsilon$  is small enough,  $N_j^1 \setminus N_\epsilon^3 \subset B(x, r/2)$ . Then  $B(x, r/2) \cap N_j^2 = \emptyset$ , and (32) is not satisfied.

Summarizing, condition (32) is satisfied if  $d(x, y) < \tau$  and not satisfied if  $d(x, y) > \tau$ . As  $d(x, y) < T$ , this yields the claim.  $\square$

Next we show that by using the processed time reversal iteration, we can test if condition (32) is valid. To this end, denote

$$\begin{aligned} \tilde{N}_1(j, \epsilon) &= N_j^1 \cup N_\epsilon^3 & \tilde{B}_1(j, \epsilon) &= B_j^1 \cup B_\epsilon^3 \\ \tilde{N}_2(j, \epsilon) &= N_j^2 \cup N_\epsilon^3 & \tilde{B}_2(j, \epsilon) &= B_j^2 \cup B_\epsilon^3 \\ \tilde{N}_3(j, \epsilon) &= N_j^1 \cup N_j^2 \cup N_\epsilon^3 & \tilde{B}_3(j, \epsilon) &= B_j^1 \cup B_j^2 \cup B_\epsilon^3 \\ \tilde{N}_4(j, \epsilon) &= N_\epsilon^3 & \tilde{B}_4(j, \epsilon) &= B_\epsilon^3. \end{aligned}$$

Let  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$ . Using the processed time reversed iteration on time interval  $[0, 2T]$  with projectors  $P_B$  corresponding to  $B = \tilde{B}_k(j, \epsilon)$ ,  $k = 1, 2, 3, 4$  and starting point  $f$ , we obtain functions  $h_n(\alpha; \epsilon, j, k) \in L^2(\partial M \times [0, 2T])$ . Using them, define

$$p_n(\alpha, j, \epsilon) = h_n(\alpha; \epsilon, j, 1) + h_n(\alpha; \epsilon, j, 2) - h_n(\alpha; \epsilon, j, 3) - h_n(\alpha; \epsilon, j, 4). \quad (33)$$

**Lemma 10** *The condition (32) is satisfied if and only if there exists an  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$  such that for any  $j \in \mathbb{Z}_+$  and  $\epsilon > 0$  the functions  $p_n(\alpha, j, \epsilon)$  defined in formula (33) satisfy*

$$\lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} \langle K_{2T} p_n(\alpha, j, \epsilon), p_n(\alpha, j, \epsilon) \rangle \neq 0. \quad (34)$$

**Proof.** The functions  $h_n(\alpha; \epsilon, j, k)$  defined by processed time reversal iteration satisfy

$$\chi_{\tilde{N}_k(j, \epsilon)} u^f(T) = \lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} u^{h_n(\alpha; \epsilon, j, k)}(T), \quad k = 1, 2, 3, 4.$$

A simple computation gives us

$$\chi_{I(j, \epsilon)}(x) = \chi_{\tilde{N}_1(j, \epsilon)}(x) + \chi_{\tilde{N}_2(j, \epsilon)}(x) - \chi_{\tilde{N}_3(j, \epsilon)}(x) - \chi_{\tilde{N}_4(j, \epsilon)}(x) \quad (35)$$

for all  $x \in M$ . Therefore, using (33) we see that in  $L^2(M)$

$$\chi_{I(j, \epsilon)} u^f(T) = \lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} u^{p_n(\alpha, j, \epsilon)}(T).$$

By Theorem 5 we see that the functions  $u^f(T)$  with  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$  are smooth functions that form a dense set in  $L^2(M(\partial M, T))$ . Thus condition (32) is satisfied if and only if there exists an  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$  such that for any  $j \in \mathbb{Z}_+$  and  $\epsilon > 0$

$$\begin{aligned} & \langle \chi_{I(j, \epsilon)} u^f(T), u^f(T) \rangle_{L^2(M)} \\ &= \lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} \langle K_{2T} p_n(\alpha, j, \epsilon), p_n(\alpha, j, \epsilon) \rangle_{L^2(\partial M \times [0, 2T])} \neq 0. \end{aligned}$$

This proves the claim.  $\square$

We note that if condition (34) is satisfied for some  $f \in C_0^\infty(\partial M \times \mathbb{R}_+)$ , it is satisfied for all such  $f$  in an open and dense set.

Lemmata 9 and 10 give an algorithm for the determination of  $d(y, x)$  by using processed time reversal iteration

$$\begin{aligned} d(y, x) = \inf \{ \tau \in [0, T] : & \text{there is an } f \in C_0^\infty(\partial M \times \mathbb{R}_+) \\ & \text{such that (34) holds for all } j \in \mathbb{Z}_+ \text{ and } \epsilon > 0 \}. \end{aligned} \quad (36)$$

Summarizing, we have proven:

**Theorem 11** *Assume we are given  $\partial M$  and the response operator  $\Lambda$ . Let  $z, y \in \partial M$ ,  $T_1 \leq \tau(z)$ . Then using the algorithm (36) we can compute  $d(x, y)$  for  $x = \gamma_{z, \nu}(T_1)$ .*

Let us consider consequences of the result above. To this end, we define the set of the boundary distance functions. For each  $x \in M$ , the corresponding boundary distance function,  $r_x \in C(\partial M)$  is given by

$$r_x : \partial M \rightarrow \mathbb{R}_+, \quad r_x(z) = d(x, z), \quad z \in \partial M.$$

In fact,  $r_x \in \text{Lip}(\partial M)$  with the Lipschitz constant equal to one. The boundary distance functions define *the boundary distance map*  $\mathcal{R} : M \rightarrow C(\partial M)$ ,  $\mathcal{R}(x) = r_x$ , which is continuous and injective (see [26]). Denote by

$$\mathcal{R}(M) = \{r_x \in C(\partial M) : x \in M\},$$

the image of  $\mathcal{R}$ . It is known (see [26, 27]) that, given the set  $\mathcal{R}(M) \subset C(\partial M)$  we can endow it, in a constructive way, with a differentiable structure and a metric tensor  $\tilde{g}$ , so that  $(\mathcal{R}(M), \tilde{g})$  becomes a manifold that is isometric to  $(M, g)$ ,

$$(\mathcal{R}(M), \tilde{g}) \cong (M, g).$$

See [27]. A stable construction procedure of  $(M, g)$  as a metric space from the set  $\mathcal{R}(M), \tilde{g}$  and Hölder type stability estimates are give in [24].

**Example** By the triangle inequality,

$$\|r_x - r_y\|_\infty \leq d(x, y). \quad (37)$$

We consider in this example the case when  $(M, g)$  is a compact manifold such that all points  $x, y \in M$  can be joined with a unique shortest geodesic. This implies that, for any  $x, y \in M$ , the shortest geodesic  $\gamma([0, s])$  from  $x$  to  $y$ , parametrised along arc length, can be continued to a maximal geodesic  $\gamma([0, L])$  that hits the boundary at a point  $z = \gamma(L) \in \partial M$ . Then

$$|r_x(z) - r_y(z)| = d(x, y)$$

implying equality in (37). Thus in the case when all geodesics between arbitrary points  $x, y \in M$  are unique, the manifold  $(M, g)$  is isometric to the manifold  $\mathcal{R}(M)$  with the distance function inherited from  $C(\partial M)$ . In the general case the construction of the metric is more elaborate.

By Theorem 11 we can compute for all  $x = \gamma_{z, \nu}(T)$  with  $T \leq \tau(z)$  the corresponding boundary distance function  $r_x$ . Since all points  $x \in M$  can be represented in this form (see e.g. [15]) we can find the set  $\mathcal{R}(M)$  that can be endowed with a manifold structure isometric to the original manifold  $(M, g)$ . We have thus proven the following result:

**Corollary 12** *Assume we are given  $\partial M$  and the response operator  $\Lambda$ . Then using the processed time reversal iteration we can find the manifold  $(M, g)$  upto an isometry.*

Corollary 12 can also be formulated by saying that we can find the metric tensor  $g$  in local coordinates. For example, for any point  $x_0 \in M$  there are  $z_j \in \partial M$ ,  $j = 1, 2, \dots, m$ , such that  $x \mapsto (X^1(x), \dots, X^m(x))$  with  $X^j(x) := d(x, z_j)$  define local coordinates near  $x_0$ . In these coordinates the distance functions  $x \mapsto d(x, z)$ ,  $z \in \partial M$  determine the metric tensor. For details of this construction, see [26].

Next we prove Corollary 3.

**Proof of Corollary 3.** As the boundary data  $\partial M$  and  $\Lambda$  determine  $(M, g)$  upto isometry, we can apply formula (31) to find the function  $C_0(x)$ ,  $x \in M$  in Corollary 2 in any local coordinates of  $(M, g)$ . Thus in local coordinates we can find the values of waves  $u^f(x, t)$  for all  $x \in M$ ,  $t > 0$ . By Tataru's controllability theorem, the waves  $u^f(x, T)$  with fixed  $T > 0$  form a dense set in  $L^2(M(\partial M, T))$ . In this dense set we can find in local coordinates the distributions

$$Au^f(x, t) = -\partial_t^2 u^f(x, t) = -u^{f''}(x, t),$$

implying that we can find values  $Aw$  for all  $w \in L^2(M(\partial M, T))$ . As  $T$  is arbitrary, we find  $Aw$  for all  $w \in L^2(M)$ .

The case when  $M \subset \mathbb{R}^m$ ,  $m \geq 2$  and  $A$  is of the form (6), we use the fact that a manifold with conformally Euclidian metric is embedded into  $\mathbb{R}^m$  in a unique way when  $\partial M$  is fixed. Thus we find  $c(x)$  when a metric  $g$  is given, and as above we also find an operator  $A_\kappa$  that is equivalent to  $A$  through a gauge transformation. As there is a unique operator of the form (6) in the orbit of the the gauge transformations  $A \rightarrow A_\kappa$  the claim follows (see [26]).  $\square$

## 6. Iteration when measurements have errors

Let  $(\Omega, \Sigma, \mathbb{P})$  be a complete probability space.

Assume the measurements have a random noise, that is, measurements give us with an input  $f$  the function  $\Lambda_{2T}f + \epsilon$ , where  $\epsilon$  is random Gaussian noise that has values in  $L^2(\partial M \times (0, T))$ . Assume that  $\mathbb{E}\epsilon = 0$  and denote the covariance operator of  $\epsilon$  by  $C_\epsilon$ . Note that  $C_\epsilon$  is a compact operator on  $L^2(\partial M \times (0, T))$  (see [12]) and thus the standard white noise on  $\partial M \times (0, T)$  does not satisfy our assumptions.

Assume that the noise is independent of previous measurements each time when we do a new measurement. When the noise is added to the processed time reversal iteration, we come to the iteration of the form

$$\tilde{h}_{n+1} = S\tilde{h}_n + p + N_n,$$

where  $N_n = PJ\epsilon_n^1 - PR\epsilon_n^2$  where  $\epsilon_n^1$  and  $\epsilon_n^2$  are random variables having the same distribution as  $\epsilon$ . Thus the  $N_n$  are independent identically distributed Gaussian random variables satisfying  $\mathbb{E}N_n = 0$  and having covariance operator  $C_N = PJC_\epsilon J^*P^* + PRC_\epsilon R^*P^*$ . Let us consider the averaged results of iterations

$$\tilde{h}_K^{ave} = \frac{1}{K} \sum_{n=1}^K \tilde{h}_n. \quad (38)$$

Then

$$\begin{aligned} \frac{1}{K} \sum_{n=1}^K \tilde{h}_n &= \frac{1}{K} \sum_{n=1}^K h_n + \frac{1}{K} \sum_{n=1}^K \sum_{m=1}^n (n-m+1) S^{n-m} N_n \\ &= \left( \frac{1}{K} \sum_{n=1}^K h_n \right) + \left( \frac{1}{K} \sum_{n=1}^K (I-S)^{-2} N_n \right) \\ &\quad + \left( \frac{1}{K} \sum_{n=1}^K (-(I-S)^{-2} S^{n+2} - (n+2)(I-S)^{-1} S^{n+1}) N_n \right) \\ &= H_K^1 + H_K^2 + H_K^3, \end{aligned}$$

where  $h_n$  are the results of the processed time reversal iteration without noise.

Above, the deterministic term  $H_K^1$  converges to the same limit  $\lim_{n \rightarrow \infty} \tilde{h}_n = h(\alpha)$  as the processed time reversal iteration without noise. Now consider  $H_K^2$  and  $H_K^3$  as random variables in  $L^2(\partial M \times (0, T))$ . They can also be viewed as random fields on  $\partial M \times (0, T)$ , see [37]. By the law of large numbers in infinite dimensional spaces [20], we see that

$$\lim_{K \rightarrow \infty} \|H_K^2\|_{L^2(\partial M \times (0, T))} = 0. \quad (39)$$

As  $\|S\|_{L(L^2(\partial M))} \leq \frac{1}{2}$ , the last term  $H_K^3$  satisfies also an estimate analogous to (39). Thus the averaged processed time reversal iteration with noise converges to the same limit as processed time reversal iteration without noise, that is,

$$\lim_{K \rightarrow \infty} \tilde{h}_K^{ave} = h(\alpha) \quad \text{in } L^1(\Omega; L^2(\partial M \times (0, T))).$$

**Acknowledgements:** The research was partially supported by the TEKES project ‘‘MASIT03 – Inverse Problems and Reliability of Models’’.

## References

- [1] M. Anderson, A. Katsuda, Y. Kurylev, M. Lassas, M. Taylor: Boundary regularity for the Ricci equation, Geometric Convergence, and Gel'fand's Inverse Boundary Problem, *Inventiones Mathematicae* 158 (2004), 261–321.
- [2] G. Aubert, P. Kornprobst: Mathematical problems in image processing. Partial differential equations and the calculus of variations. Applied Mathematical Sciences, 147. Springer, New York, 2006. xxxii + 377 pp.
- [3] C. Bardos: A mathematical and deterministic analysis of the time-reversal mirror. *Inside out: inverse problems and applications*, 381–400, Math. Sci. Res. Inst. Publ., 47, Cambridge Univ. Press, Cambridge, 2003.
- [4] C. Bardos, M. Fink: Mathematical foundations of the time reversal mirror. *Asymptot. Anal.* 29 (2002), no. 2, 157–182.
- [5] G. Bal, G. Papanicolaou, L. Ryzhik: Self-averaging in time reversal for the parabolic wave equation, *Stochastics and Dynamics* 2 (2002), 507–531
- [6] G. Bal, O. Pinaud: Time-reversal-based detection in random media. *Inverse Problems* 21 (2005), 1593–1619.
- [7] G. Bal, L. Ryzhik: Time reversal for classical waves in random media. *C. R. Acad. Sci. Paris Sér. I Math.* 333 (2001), 1041–1046.
- [8] G. Bal, R. Verástegui: Time reversal in changing environments. *Multi-scale Model. Simul.* 2 (2004), 639–661
- [9] M. Belishev: An approach to multidimensional inverse problems for the wave equation. (Russian) *Dokl. Akad. Nauk SSSR* 297 (1987), no. 3, 524–527.
- [10] M. Belishev: Boundary control in reconstruction of manifolds and metrics (the BC method). *Inverse Problems* 13 (1997), no. 5, R1–R45.
- [11] M. Belishev, Y. Kurylev: To the reconstruction of a Riemannian manifold via its spectral data (BC-method). *Comm. Partial Differential Equations* 17 (1992), 767–804.

- [12] V.I. Bogachev: Gaussian measures. Mathematical Surveys and Monographs, 62. American Mathematical Society, Providence, RI, 1998. xii + 433 pp.
- [13] L. Borcea, G. Papanicolaou, C. Tsogka, J. Berryman: Imaging and time reversal in random media. *Inverse Problems* 18 (2002), 1247–1279.
- [14] L. Borcea, G. Papanicolaou, C. Tsogka: Theory and applications of time reversal and interferometric imaging. Special section on imaging. *Inverse Problems* 19 (2003), S139–S164.
- [15] I. Chavel: Riemannian geometry. A modern introduction. Cambridge University Press, Cambridge, 2006. xvi + 471 pp.
- [16] M. Cheney, D. Isaacson, M. Lassas: Optimal acoustic measurements. *SIAM J. Appl. Math.* 61 (2001), no. 5, 1628–1647.
- [17] M. Fink: Time reversal mirrors. *J. Phys. D: Appl. Phys.* 26 (1993), 1333–1350.
- [18] M. Fink: Acoustic Time-Reversal Mirrors. *Imaging of Complex Media with Acoustic and Seismic Waves*, 17–43. Topics in Applied Physics, 84. Springer, 2002.
- [19] M. Fink, D. Cassereau, A. Derode, C. Prada, P. Roux, M. Tanter, J.-L. Thomas, F. Wu: Time-reversed acoustics. *Rep. Prog. Phys.* 63 (2000), 1933–1995.
- [20] J. Hoffmann-Jorgensen, G. Pisier: The Law of Large Numbers and the Central Limit Theorem in Banach Spaces. *The Annals of Probability* 4 (1976), 587–599.
- [21] L. Hörmander: The analysis of linear partial differential operators. III. Pseudodifferential operators. Springer-Verlag, 1985. viii + 525 pp.
- [22] D. Isaacson: Distinguishability of conductivities by electric current computed tomography. *IEEE Trans. on Medical Imaging* MI-5 (1986), 92–95.
- [23] B.L.G. Jonsson, M. Gustafsson, V.H. Weston, M.V. de Hoop: Retrofocusing of acoustic wave fields by iterated time reversal. *SIAM J. Appl. Math.* 64 (2004), no. 6, 1954–1986.

- [24] A. Katsuda, Y. Kurylev, M. Lassas: Stability of boundary distance representation and reconstruction of Riemannian manifolds, *Inverse Problems and Imaging* 1 (2007), 135–157.
- [25] A. Katchalov, Y. Kurylev: Multidimensional inverse problem with incomplete boundary spectral data. *Comm. Part. Diff. Equations* 23 (1998), no. 1–2, 55–95.
- [26] A. Katchalov, Y. Kurylev, M. Lassas: Inverse boundary spectral problems. Chapman & Hall/CRC, 2001. xx + 290 pp.
- [27] A. Katchalov, Y. Kurylev, M. Lassas: Energy measurements and equivalence of boundary data for inverse problems on non-compact manifolds. *Geometric methods in inverse problems and PDE control* (eds. C. Croke, I. Lasiecka, G. Uhlmann, M. Vogelius), 183–213, IMA volumes in Mathematics and Applications, 137, Springer, New York, 2004.
- [28] A. Katchalov, Y. Kurylev, M. Lassas, N. Mandache: Equivalence of time-domain inverse problems and boundary spectral problem, *Inverse problems* 20 (2004), 419–436.
- [29] M. Klibanov, A. Timonov: On the mathematical treatment of time reversal. *Inverse Problems* 19 (2003), 1299–1318.
- [30] Y. Kurylev, M. Lassas: Hyperbolic inverse problem with data on a part of the boundary. *Differential Equations and Mathematical Physics (Birmingham, AL, 1999)*, AMS/IP Stud. Adv. Math., 16 (2000), 259–272, Amer. Math. Soc., Providence, 2000.
- [31] Y. Kurylev, M. Lassas: Hyperbolic inverse boundary-value problem and time-continuation of the non-stationary Dirichlet-to-Neumann map. *Proc. Roy. Soc. Edinburgh Sect. A* 132 (2002), no. 4, 931–949.
- [32] I. Lasiecka, R. Triggiani: Sharp regularity theory for second order hyperbolic equations of Neumann type. I.  $L_2$  nonhomogeneous data. *Ann. Mat. Pura Appl. (4)* 157 (1990), 285–367.
- [33] I. Lasiecka, R. Triggiani: Regularity theory of hyperbolic equations with nonhomogeneous Neumann boundary conditions. II. General boundary data. *J. Differential Equations* 94 (1991), 112–164.
- [34] T.D. Mast, A.I. Nachman, R.C. Waag: Focusing and imaging using eigenfunctions of the scattering operator. *J. Acoust. Soc. Am.* 102 (1997) 715–725.

- [35] C. Prada, M. Fink: Eigenmodes of the time reversal operator: A solution to selective focusing in multiple-target media. *Wave Motion* 20 (1994), 151–163.
- [36] C. Prada, J.-L. Thomas, M. Fink: The iterative time reversal process: Analysis of the convergence. *J. Acoust. Soc. Am.* 97 (1995), 62–71.
- [37] Yu. Rozanov: Markov random fields. Springer-Verlag, New York-Berlin, 1982. ix + 201 pp.
- [38] D. Tataru: Unique continuation for solutions to PDEs, between Hörmander’s theorem and Holmgren’s theorem. *Comm. Part. Diff. Equations* 20 (1995), 855–884.
- [39] D. Tataru: On the regularity of boundary traces for the wave equation. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* 26 (1998), 185–206.
- [40] D. Tataru: Unique continuation for operators with partially analytic coefficients. *J. Math. Pures Appl.* 78 (1999), 505–521.

Kenrick Bingham and Matti Lassas  
 Institute of Mathematics  
 P.O.Box 1100  
 02015 Helsinki University of Technology  
 Finland

Yaroslav Kurylev  
 Department of Mathematical Sciences  
 Loughborough University  
 Loughborough  
 Leicestershire  
 LE11 3TU  
 UK

Samuli Siltanen  
 Institute of Mathematics  
 Tampere University of Technology  
 P.O.Box 553  
 33101 Tampere  
 Finland